

After a Concussion:

RETURN-TO-SPORT STRATEGY



Parachute
Concussion Series

A concussion is a serious injury, but you can recover fully if your brain is given enough time to rest and recuperate.

Returning to normal activities, including sport participation, is a step-wise process that requires patience, attention, and caution.

In the Return-to-Sport Strategy:

- ▶ Each stage is at least 24 hours.
- ▶ Move on to the next stage when activities are tolerated without new or worsening symptoms.
- ▶ If any symptoms worsen, stop and go back to the previous stage for at least 24 hours.
- ▶ If symptoms return after medical clearance, follow up with a doctor for re-assessment.

Stage 1: Symptom-limiting activities

After an initial short period of rest of 24 to 48 hours, light cognitive and physical activity can begin, as long as these don't worsen symptoms. You can start with daily activities like moving around the house, simple chores, and gradually introducing school and work activities at home.

Stage 2: Light aerobic activity

Light exercise such as walking or stationary cycling, for 10 to 15 minutes. The duration and intensity of the aerobic exercise can be gradually increased over time if symptoms don't worsen and no new symptoms appear during the exercise or in the hours that follow. No resistance training or other heavy lifting.

Stage 3: Individual sport-specific exercise with no contact

Activities such as skating, running, or throwing can begin for 20 to 30 minutes. There should be no body contact or other jarring motions, such as high-speed stops or hitting a ball with a bat. No resistance training.

Stage 4: Begin training drills with no contact

Add in more challenging drills like passing drills. There should be no impact activities (no checking, no heading the ball, etc.). Start to add in progressive resistance training.

Stage 5: Full contact practice following clearance by a doctor.

Stage 6: Return to Sport

Full game play or competition.



Parachute is Canada's
leading national charity
dedicated to injury prevention.



For concussion info on the go, download the app



Appendix C-2

Sample Tool to Identify a Suspected Concussion¹

This sample tool is a quick reference, to be completed by teachers, to help identify a suspected concussion and to communicate this information to parent/guardian.

Identification of Suspected Concussion

Following a blow to the head, face or neck, or a blow to the body that transmits a force to the head, a concussion must be suspected in the presence of **any one or more** of the signs or symptoms outlined in the chart below and/or the failure of the Quick Memory Function Assessment.

First, assess the danger to the student and the rescuer, and then check airway, breathing and circulation.

1. Check appropriate box

An incident occurred involving _____ (student name) on _____ (date).
He/she was observed for signs and symptoms of a concussion.

- No signs or symptoms described below were noted at the time. **Note:** *Continued monitoring of the student is important as signs and symptoms of a concussion may appear hours or days later (refer to #4 below).*
- The following signs were observed or symptoms reported:

Signs and Symptoms of Suspected Concussion	
Possible Signs Observed <i>A sign is something that is observed by another person (e.g., parent/guardian, teacher, coach, supervisor, peer).</i>	Possible Symptoms Reported <i>A symptom is something the student will feel/report.</i>
<p>Physical</p> <ul style="list-style-type: none"> <input type="checkbox"/> vomiting <input type="checkbox"/> slowed reaction time <input type="checkbox"/> amnesia <input type="checkbox"/> blank stare/glassy-eyed/dazed or vacant look <input type="checkbox"/> loss of consciousness or lack of responsiveness (<i>call 911 immediately</i>) <input type="checkbox"/> lying motionless on the ground or slow to get up <input type="checkbox"/> seizure or convulsion (<i>call 911 immediately</i>) <input type="checkbox"/> grabbing or clutching of head <p>Cognitive</p> <ul style="list-style-type: none"> <input type="checkbox"/> difficulty concentrating <input type="checkbox"/> easily distracted <input type="checkbox"/> general confusion <input type="checkbox"/> cannot remember things that happened before and after the injury (<i>see Quick Memory Function Assessment on page 2</i>) <input type="checkbox"/> does not know time, date, place, class, type of activity in which he/she was participating <input type="checkbox"/> slowed reaction time (e.g., answering questions or following directions) <p>Emotional/Behavioural</p> <ul style="list-style-type: none"> <input type="checkbox"/> strange or inappropriate emotions (e.g., laughing, crying, getting angry easily) <p>Other</p> <ul style="list-style-type: none"> <input type="checkbox"/> _____ 	<p>Physical</p> <ul style="list-style-type: none"> <input type="checkbox"/> headache <input type="checkbox"/> pressure in head <input type="checkbox"/> neck pain <input type="checkbox"/> feeling off/not right <input type="checkbox"/> ringing in the ears <input type="checkbox"/> seeing double or blurry/loss of vision <input type="checkbox"/> seeing stars, flashing lights <input type="checkbox"/> pain at physical site of injury <input type="checkbox"/> nausea/stomach ache/pain <input type="checkbox"/> balance problems or dizziness <input type="checkbox"/> fatigue or feeling tired <input type="checkbox"/> sensitivity to light or noise <p>Cognitive</p> <ul style="list-style-type: none"> <input type="checkbox"/> difficulty concentrating or remembering <input type="checkbox"/> slowed down, fatigue or low energy <input type="checkbox"/> dazed or in a fog <p>Emotional/Behavioural</p> <ul style="list-style-type: none"> <input type="checkbox"/> irritable, sad, more emotional than usual <input type="checkbox"/> nervous, anxious, depressed <p>Other</p> <ul style="list-style-type: none"> <input type="checkbox"/> _____

CONCUSSION RECOGNITION TOOL 5 ©

To help identify concussion in children, adolescents and adults



Supported by



RECOGNISE & REMOVE

Head impacts can be associated with serious and potentially fatal brain injuries. The Concussion Recognition Tool 5 (CRT5) is to be used for the identification of suspected concussion. It is not designed to diagnose concussion.

STEP 1: RED FLAGS —CALL AN AMBULANCE

If there is concern after an injury including whether ANY of the following signs are observed or complaints are reported then the player should be safely and immediately removed from play/games and they should not be assessed. Healthcare professional is available, call an ambulance for urgent medical assessment:

- Neck pain or tenderness
- Double vision
- Weakness or tingling/ burning in arms or legs
- Severe or increasing headache
- Seizure or convulsion
- Loss of consciousness
- Deteriorating conscious state
- Vomiting
- Increasingly restless, agitated or combative

Remember:

- In all cases, the basic principles of first aid (danger, response, airway, breathing, circulation) should be followed.
- Assessment for a spinal cord injury is critical.
- Do not attempt to move the player (other than required for airway support) unless trained to do so.
- Do not remove a helmet or any other equipment unless trained to do so safely.

If there are no Red Flags, identification of possible concussion should proceed to the following steps:

STEP 2: OBSERVABLE SIGNS

Visual clues that suggest possible concussion include:

- Lying motionless on the playing surface
- Slow to get up after a direct or indirect hit to the head
- Disorientation or confusion, or an inability to respond appropriately to questions
- Balance, gait difficulties, motor incoordination, stumbling, slow laboured movements
- Blank or vacant look
- Facial injury after head trauma

© Concussion in Sport Group 2017

STEP 3: SYMPTOMS

- Headache
- "Pressure in head"
- Balance problems
- Nausea or vomiting
- Drowsiness
- Dizziness
- Blurred vision
- Sensitivity to light
- Sensitivity to noise
- Fatigue or low energy
- "Don't feel right"
- More emotional
- More irritable
- Sadness
- Nervous or anxious
- Neck Pain
- Difficulty concentrating
- Difficulty remembering
- Feeling slowed down
- Feeling like "in a fog"

STEP 4: MEMORY ASSESSMENT

(IN ATHLETES OLDER THAN 12 YEARS)

- Failure to answer any of these questions (modified appropriately for each sport) correctly may suggest a concussion:
- "What venue are we at today?"
 - "Which half is it now?"
 - "Who scored last in this game?"
 - "What team did you play last week/game?"
 - "Did your team win the last game?"

Athletes with suspected concussion should:

- Not be left alone initially (at least for the first 1-2 hours).
- Not drink alcohol.
- Not use recreational/ prescription drugs.
- Not be sent home by themselves. They need to be with a responsible adult.
- Not drive a motor vehicle until cleared to do so by a healthcare professional.

The CRT5 may be freely copied in its current form for distribution to individuals, teams, groups and organisations. Any revision and any reproduction in a digital form requires approval by the Concussion in Sport Group. It should not be altered in any way, rebranded or sold for commercial gain.

ANY ATHLETE WITH A SUSPECTED CONCUSSION SHOULD BE IMMEDIATELY REMOVED FROM PRACTICE OR PLAY AND SHOULD NOT RETURN TO ACTIVITY UNTIL ASSESSED MEDICALLY, EVEN IF THE SYMPTOM IS RESOLVE

© Concussion in Sport Group 2017

Facility Safety Checklist

Facility: _____

Date: _____

Inspected by: _____

Item	Adequate	Inadequate	Corrective Measures*	Observations
Playing surface and installations				
Dressing Room				
Team Equipment				
Individual Equipment				

*Corrections: 1-add 2-replace 3-modify 4-discard 5-clean 6-repair 7-check